

## **Important Safety Information about Taclonex Scalp<sup>®</sup> Topical Suspension**

Taclonex Scalp<sup>®</sup> is for the treatment of moderate to severe psoriasis vulgaris on the scalp in adults, and should be applied to affected areas once daily for 2 weeks or until cleared. You may use it for up to 8 weeks or as prescribed by your doctor, and you should not use more than 100 grams per week.

**FOR TOPICAL USE ONLY.** Do not use on your face, under your arms or on your groin. **Do not swallow.**

You should not use Taclonex Scalp<sup>®</sup> if you have a history of allergy to any of its ingredients; thin skin (atrophy) at the treatment site; severe kidney or liver disease; higher than normal levels of calcium in your blood; or erythrodermic, exfoliative or pustular psoriasis.

Serious side effects may include too much calcium in your blood, or adrenal gland problems if you use too much, use it for too long, or use it with other topical medicines that contain corticosteroids or calcipotriene. The most common side effects are itching, inflamed hair pores (folliculitis), skin pain or irritation, and skin burning at the site of application.

If you are pregnant or plan to become pregnant while using Taclonex Scalp<sup>®</sup>, talk to your doctor before using this medication. You should not bandage, tightly cover or wrap the treated skin area. Other products containing calcipotriene or a corticosteroid should not be used with Taclonex Scalp<sup>®</sup> without first talking to your doctor. Use sunscreen or wear a hat if you plan to be in sunlight for a long period of time.

Please refer to the Patient Information for additional important information.